

# CLINICAL WORKER WELLBEING: SAFETY IN CLINICIAN WORKSPACES

## QUANTITATIVE RESEARCH

**300+**

interviews with nurses and clinicians across the U.S. and Canada


## QUALITATIVE RESEARCH

**2,000 HOURS**

observing interactions around patient seating


Clinicians are concerned about, and vulnerable to, work-related injuries.

### CLINICIAN INJURY CONCERN




### CLINICIAN INJURY EXPERIENCE

**48%** of clinicians surveyed have been impacted in some way by clinical injuries.



Clinicians may be more likely to get hurt because they are asked to do more than ever before.

### CLINICIANS RATE DIRECT PATIENT CARE ACTIVITIES AS HIGH-RISK



### PATIENT TRANSFERS ARE ESPECIALLY RISKY

**1 IN 3** clinicians surveyed has experienced an injury related to patient transfers.




### MOST COMMON INJURIES

Clinicians	Patients
Back Injury	Falling
Neck Injury	Bruising
Pain	Pain

Managing risk: more can be done to support nurses and clinicians.

### CLINICIANS ARE ASKING FOR MORE SUPPORT



### CLINICIANS DESIRE MORE SUPPORTIVE SPACES


#### Five Key Insights

- High risk of injury on the job
- Bed-to-chair transfer causes most injuries
- Clinicians try to personally maintain their health and safety on the job
- Clinicians desire one common change in their work environments
- Supporting a selfless profession

Empath recliner: clinicians believe it can help


### CLINICIANS ARE INTERESTED IN EMPATH

**68%** indicate an interest in Empath as a support in their work environment.



### EMPATH VERSUS OTHER RECLINERS

**81%** believe that Empath is superior to other recliners.



### CLINICIANS AGREE THAT EMPATH ENHANCES EXPERIENCE

